Adult Immunization Schedule

Adolescents and Young Adults, 18 to 25 Years:

VACCINE	DOSE/ROUTE	INITIAL SERIES	BOOSTER
Tetanus and Diphtheria	0.5 cc IM	3 doses (0, 1 mos., 6-12 mos. After 2nd dose)	Every 10 years
MMR	0.5 cc SQ	2 doses (0, 1 mos.)	Not required
Influenza	0.5 cc IM	N/A	Annually
Varicella	0.5 cc SQ	2 doses (0, 1-2 mos. After 2nd dose)	See below *

Adults, 25 to 64 Years:

VACCINE	DOSE/ROUTE	INITIAL SERIES	BOOSTER
Tetanus and Diphtheria	0.5 cc IM	3 doses (0, 1 mos., 6-12 mon after 2nd dose)	Every 10 years
MMR	0.5cc SQ	2 doses (0, 1 mos.)	See below **
Influenza	0.5 cc IM	N/A	Annually
Pneumococcal	0.5cc IM or SQ	See below #	See below ##
Varicella	0.5 cc SQ	2 doses (0, 1-2 mos. After 2nd dose)	See below *

Adults, 65 Years and Older:

VACCINE	DOSE/ROUTE	INITIAL SERIES	BOOSTER
Tetanus and Diphtheria	0.5cc IM	3 doses	Every 10 years
Influenza	0.5cc IM	N/A	Annually
Pneumococcal	0.5cc SQ or IM	1 dose	See below ##

* People with a reliable history of chickenpox such as self or parental report of disease can be considered to be immune
** MMR is only required for adults born before 1957
# May be required for adults 50-64 years of age with risk factors
Every 5 years, only for elder adults at highest risk